

Nut Trees (Not Bearing)

Macadamia, Pecan, Pistachio, Brazil, Cashew

| GROWTH STAGE | PRODUCTS | Average Dosage / 100Lt Water | Maximum Dosage / Hectare | Water Volume / Hectare | COMMENTS |
|-------------------------|---|------------------------------|--------------------------|------------------------|--|
| Non Bearing Fruit trees | Amino Gluconate Zn | 160ml | 8Lt | 2000 - 3500Lt | Apply at monthly intervals, between August and March |
| New Flush | Man-O-CuZin | 500ml | 3Lt | | Apply twice during this growth period |
| Additional Note | Use Maxi-Kelp as a growth booster at a dosage of 350-500ml/100Lt water at a maximum of 8-15Lt/ha. First application after leaves have emerged. Repeat 4-6 times | | | | |

Nut Trees (Bearing)

Macadamia, Pecan, Pistachio, Brazil, Cashew

| GROWTH STAGE | PRODUCTS | Average Dosage / 100Lt Water | Maximum Dosage / Hectare | Water Volume / Hectare | COMMENTS |
|------------------------|--|------------------------------|--------------------------|------------------------|--|
| Post Harvest | Amino Gluconate Zn | 160ml | 8Lt | 2000 - 3500Lt | Apply at monthly intervals, between August and March |
| New Flush | 1:0:0 (30) LB Urea | 1400ml | 40Lt | | Apply once during this growth period |
| | ZinCoBor | 275ml | 4Lt | | |
| Flowering | Man-O-CuZin | 500ml | 3Lt | | Apply twice during this growth period |
| | Amino Gluconate K | 600ml | 4Lt | | |
| Fruiting | Amino Gluconate Ca | 200ml | 6Lt | | Apply three times during this growth period |
| | Amino Gluconate K | 600ml | 4Lt | | |
| Ripening | Amino Gluconate Ca | 200ml | 6Lt | | Apply once during this growth stage |
| | Amino Gluconate K | 600ml | 4Lt | | |
| | Amino Gluconate Zn | 80ml | 1Lt | | |
| Additional Note | Use Maxi-Kelp as a growth and fruit booster at a dosage of 350-500ml/100Lt water at a maximum of 8-15Lt/ha. First application after leaves have emerged. Repeat 4-6 times. | | | | |